

ERRATUM

On page 296 of an article by Rosenberg et al (*Biol Psychiatry* 2000;48:294–300), the clinical inventory data published in Table 2 contained minor errors. The correct

data appears below. The data concerning brain volumes in the original Table 2 contained no errors.

Table 2. Clinical Assessment Data from Psychotropic Medication-Naïve Pediatric Patients with Obsessive-Compulsive Disorder (OCD) before and after Cognitive Behavioral Therapy (CBT)

Instrument	Subjects ^a							
	Patients with OCD before treatment (mean ± SD)		Patients with OCD after CBT (mean ± SD)		<i>t</i> ^b		<i>p</i>	
	Incorrect	Correct	Incorrect	Correct	Incorrect	Correct	Incorrect	Correct
Total score CY-BOCS	22.45 ± 4.16	22.55 ± 5.22	10.45 ± 5.48	9.91 ± 5.47	5.41	5.10	.000	.000
Obsessive subscale score	10.82 ± 2.64	11.27 ± 2.83	4.73 ± 2.87	4.82 ± 2.89	4.72	4.60	.001	.001
Compulsive subscale score	11.64 ± 2.34	11.27 ± 2.69	5.64 ± 2.80	5.09 ± 2.81	5.24	5.23	.000	.000
HAM-A	7.91 ± 5.99	8.91 ± 4.83	2.00 ± 2.72	1.91 ± 2.70	3.18	4.42	.01	.001
HAM-D	7.09 ± 4.91	9.73 ± 5.85	2.18 ± 2.82	2.27 ± 2.83	3.33	3.75	.008	.004

CY-BOCS, children's version of the Yale-Brown Obsessive Compulsive Scale; HAM-A, Hamilton Anxiety Rating Scale; HAM-D, Hamilton Depression Rating Scale.

^aThere were 11 OCD patients studied before and after cognitive behavioral therapy.

^bPaired *t* test, *df* = 10.